

ON THE REMOVABLE SINGULARITIES FOR MEROMORPHIC MAPPINGS

E.M.CHIRKA

ABSTRACT. If E is a nonempty closed subset of the locally finite Hausdorff $(2n-2)$ -measure on an n -dimensional complex manifold Ω and all points of E are nonremovable for a meromorphic mapping of $\Omega \setminus E$ into a compact Kähler manifold, then E is a pure $(n-1)$ -dimensional complex analytic subset of Ω .

1. This paper was inspired by the following question of E.L.Stout [7]: Let E be a closed subset of the complex projective space $\mathbb{P}^n$ ($n \geq 2$) such that the Hausdorff $(2n-2)$ -measure of E (with respect to the Fubini-Study metric) is less than that of any complex hyperplane of $\mathbb{P}^n$. If it is true that E is then a removable singularity for meromorphic functions? Using natural projections of $\mathbb{P}^n$ onto hyperplanes G.Lupaciolu [5] has shown the removability of E under additional conditions on the sizes of E and a maximal ball in the complement. (The projection of $\mathbb{P}^n$ onto a hyperplane does not decrease Hausdorff measures, as it takes place in the Euclidean space.) Using an Oka–Nishino theorem [6] on pseudoconcave sets we prove here the following.

Theorem. *Let E be a closed subset of the locally finite Hausdorff $(2n-2)$ -measure on an n -dimensional complex manifold Ω and let f be a meromorphic mapping of $\Omega \setminus E$ into a complex manifold X . If X has the meromorphic extension property and E does not contain any $(n-1)$ -dimensional closed analytic subset of Ω then f is continued to a meromorphic mapping of Ω into X .*

Here we say that X has the meromorphic extension property, if any meromorphic map $\varphi : T \rightarrow X$ of the "squeezed polydisc"

$$T = (z, w) \in \mathbb{C}_z^{n-1} \times \mathbb{C}_w : |z| < r, |w| < 1 \text{ or } |z| < 1, 1-r < |w| < 1,$$

$0 < r < 1$, $n \geq 2$, extends to a meromorphic map $\tilde{\varphi} : U \rightarrow X$ of the unit polydisc $U : |z| < 1, |w| < 1$. By a recent result of S.M.Ivashkovich [3] every compact Kähler manifold X has the meromorphic extension property, so we have a lot of nice examples of such X . The case of meromorphic functions ($X = \mathbb{P}^1$) is almost trivial in the consideration: every meromorphic function in a squeezed polydisc is represented as a ratio of two holomorphic functions (see [4]) and thus it is meromorphically continued into U . So the answer on the question of Stout is

1991 *Mathematics Subject Classification.* Primary 32D15, 32D20; Secondary 32D10.

Key words and phrases. meromorphic continuation, removable singularities, pseudoconcave sets.

The final version of this paper will be submitted for publication elsewhere

positive because the mentioned E can not contain any complex analytic subset of $\mathbb{P}^n$ (by a Chow theorem such a set is algebraic, and the Hausdorff $(2n - 2)$ -measure of it is not less then the measure of a complex hyperplane (see, e.g. [C]). As pointed me E.L.Stout, this question was solved already by his student Mark Lawrence by a similar method.

2. A closed subset Σ of a complex manifold is called locally pseudoconcave if for every point $a \in \Sigma$ there exists a Stein neighbourhood V such that $V \setminus \Sigma$ is Stein.

Let E' be the set of points $a \in E$ such that f meromorphically extends into a neighbourhood of a . Then $S := E \setminus E'$ is closed. As the complement to E is locally connected in Ω , these local meromorphic continuations of f in points of E' glue together into the unique meromorphic map $f : \Omega \setminus S \rightarrow X$ (we preserve the notation f).

The proof of the Theorem is accomplished now in two steps. Firstly we prove that S is locally pseudoconcave (Lemma 1), and secondly we prove that S is complex analytic (Lemma 2).

Lemma 1. *S is locally pseudoconvex in Ω .*

Proof. Let $a \in S$, $V \ni a$ is biholomorphic to a ball in $\mathbb{C}^n$ and $\varphi : T \rightarrow V \setminus S$ is a holomorphic embedding. Then (V is Stein) φ extends to a holomorphic embedding $\tilde{\varphi} : U \rightarrow V$ (see [2]). As X has the meromorphic extension property, the meromorphic map $f \circ \varphi : T \rightarrow X$ extends to a meromorphic map of U into X and thus f meromorphically continues into the domain $\tilde{\varphi}(U) \subset V$. By the definition of S , $\tilde{\varphi}(U)$ does not intersect S , and thus we have proved that $V \setminus S$ satisfies the condition " p_7 -convexity" of Docquier–Grauert [2]. It follows that $V \setminus S$ is Stein.

Lemma 2. *Let S be a nonempty locally pseudoconcave subset of the finite Hausdorff $(2n - 2)$ -measure on an n -dimensional complex manifold Ω . Then S is a pure $(n - 1)$ -dimensional complex analytic subset of Ω .*

Proof. The statement is local, so we can assume that Ω is a domain in $\mathbb{C}^n$. Let $a \in S$ and coordinates (z, w) , $z = (z_1, \dots, z_{n-1})$ in $\mathbb{C}^n$ are chosen in such a way that $a = 0$ and the set $S \cap \{z = 0\}$ is finite or countable (it can be done obviously). Then there exists a neighbourhood $V = V' \times V_n \ni 0$ such that the projection of $S \cap V$ into V' is proper. It follows that fibres $S \cap V \cap \{z = c\}$ are finite for almost every $c \in V'$. By an Oka–Nishino theorem [6] $S \cap V$ is then a complex analytic set of pure dimension $n - 1$.

I would like to thank G.Lupaccolu for the reprint of his paper [5], which stimulated this work.

REFERENCES

1. E. M. Chirka, *Complex analytic sets*, "Nauka", Moscow, 1985 (Russian); English transl., Kluwer Acad. Publishers, Dordrecht-Boston-London, 1989.
2. F. Docquier and H. Grauert, *Levisches Problem und Rungescher Satz für Teilgebiete Steinscher Mannigfaltigkeiten*, Math. Ann. **140** (1960), 94–123..
3. S. M. Ivashkovich, *The Hartogs-type extension theorem for meromorphic maps into compact Kähler manifolds*, preprint (1990), Bochum, 1–13.
4. J. Kajiwar and E. Sakai, *Generalization of Levi–Oka's theorem concerning meromorphic functions*, Nagoya Math. J. **29** (1967), 75–84.
5. G. Lupaccolu, *Meromorphic continuation in projective space*, Bollettini U. M. I. (7) **4-A** (1990), 391–397.

6. T. Nishino, *Sur les ensembles pseudoconcaves*, J. Math. Kyoto Univ. **1-2** (1962), 225–245.
7. E. L. Stout, *Removable sets for holomorphic functions of several complex variables*, Publ. Math. **33** (1989), 345–362.

STEKLOV MATHEMATICAL INSTITUTE, VAVILOV ST. 42, MOSCOW GSP-1, 117966 RUSSIA
E-mail address: `chirka@mph.mian.su`